

OCCASIONAL PAPERS

Museum of Texas Tech University

Number 221

19 May 2003

HABITAT USE BY BIRDS AND MAMMALS ALONG THE URBAN SOUTH PLATTE RIVER IN DENVER, COLORADO

CHERI A. JONES, RONALD D. BEANE, AND ELIZABETH A. DICKERSON

INTRODUCTION

Fewer studies of vertebrates have been conducted in urban areas than on sites in more natural conditions. One urban area that has received little scientific attention is the stretch of the South Platte River passing through Denver, Denver County, Colorado. The South Platte played a key role in American history because it is located in the arid shortgrass steppe of the western United States. In this steppe characterized by extremes in temperature and precipitation, the South Platte has long provided water, shelter, food, and travel corridors for people and for wildlife.

The South Platte watershed covers more than 8,878 km² (23,000 mi²), of which 79% is in Colorado (U.S. EPA, 1999). Ute, Comanche, Arapaho, and Cheyenne used this watershed long before the arrival of settlers from Mexico and the United States. The first European settlements included fur-trading posts and forts (U.S. EPA, 1999). The earliest known European-Mexican settlement in the Denver area was the short-lived Mexican Diggings, established in 1857. Modern Den-

ver grew from the small towns of Saint Charles and Auraria, founded in 1858 near the confluence of the South Platte and Cherry Creek (Jones and Forrest, 1973; Noel, 1980; Kirkpatrick, 1998). Anthropogenic changes accompanying European settlement included the construction of canals, producing a system that presently delivers water to 29 cities and towns, 120 ditch companies, 60 reservoir companies, and 620,000 acres of irrigated farmland (U.S. EPA, 1999). As in most cities, water also provided avenues for transportation and waste disposal, leading to the establishment of railyards and industry along the river. Floods have occurred periodically, including major events in Denver in 1864 and 1965 (Meister, 1965; Brenneman, 1973; Jones and Forrest, 1973; U.S. EPA, 1999; Chapman, 2001). Human and economic losses resulting from these floods led to the establishment of reservoirs such as Chatfield (H. E. Kingery, pers. comm., 2001) and water-management projects, which have greatly altered the water flow and biotic communities of the South Platte (Johnson, 1994).

Denver County lies within the Front Range of Colorado, an area in which the human population has expanded from 2,694,000 in 1990 to 3,512,779 in 2000 (Colorado State Demographer, 2000 Census, www.dlg.oem2.state.co.us). Approximately 16 km of the South Platte pass through modern Denver, where it remains a center of activity, serving as a major focus of economic and residential development. Additionally, the river and its associated urban parks are important recreational resources for bikers, walkers, kayakers, and others (e.g., Muntz and Wuth, 1983; Hekkers, 1987; Cooper et al., 1990; Kirkpatrick, 1998; Moorhead, 1992).

In recognition of the river's importance to local residents and wildlife, Mayor William McNichols created the Platte River Development Commission in the late 1970s (Chapman, 2001). Mayor Wellington Webb assembled the South Platte River Working Group in 1994 to address improvements to the urban corridor as a major issue during his tenure (Mayor's South Platte River Commission, 1995; Instream Issues Task Force, 1996; City and County of Denver, 2000). Subsequent improvements of recreational trails (40 km, or 25 miles, along the Platte, Cherry Creek, Bear Creek, Highline Canal, and Sand Creek) resulted in one of the best greenway systems in the United States (Shuster, 1996; Anonymous, 1998; U.S. EPA, 1999). Clean-up programs, economic development, and other activities received public recognition in various publications (e.g., Miller, 1996; Anonymous, 1997, 1998; Walker, 1997; Brovsky, 1999), the 1997 American Rivers Gold Medal for Urban River Partnership, and the 1999 National Wildlife Federation Conservation Achievement Award (Alston, 1999).

However, most resource analyses of the urban corridor have concentrated on fish, hydrology, and water quality (Instream Issues Task Force, 1996; Kimbrough and Litke, 1996). Some surveys of wildlife have been conducted within the metropolitan area, but the most detailed studies of birds and mammals associated with the South Platte River have been conducted in rural Colorado (Mitchell, 1972; Fitzgerald, 1978; Olson and Knopf, 1988). Thus the potential of the urban South Platte as wildlife habitat has remained largely unexplored.

Some of the surveys along the urban South Platte include Audubon bird counts and the Urban Denver Christmas Bird Count conducted since 1988 (H. E. Kingery, pers. comm., 1999). On 1 January 2000, for the 1999-2000 Christmas Bird Count period, 56 species and 8,896 individuals were recorded along the river. Of that total, 25% were Canada geese and 25% were starlings (Kingery, 2000; please see Appendix A for taxa reported in our study), and three new species (the ruddy duck [*Oxyura jamaicensis*], mew gull [*Larus canus*], and red-shouldered hawk [*Buteo lineatus*]) were added. Thompson (1994) monitored wildlife on a 16-acre wetland-mitigation area adjacent to the river. Thirty-four avian species were reported, 70% of which were red-winged blackbirds (*Agelaius phoeniceus*), barn swallows (*Hirundo rustica*), black-billed magpies (*Pica pica*), common grackles (*Quiscalus quiscula*), mallards (*Anas platyrhynchos*), starlings, and Canada geese.

Several short-term, local surveys of mammals also were conducted. Brittan (1992) briefly surveyed small mammals in South Platte Park, capturing deer mice (*Peromyscus maniculatus*) and meadow voles (*Microtus pennsylvanicus*). Erickson (1982) documented 11 species of mammals along the South Platte Greenway: the eastern cottontail (*Sylvilagus floridanus*), fox squirrel (*Sciurus niger*), American beaver (*Castor canadensis*), western harvest mouse (*Reithrodontomys megalotis*), Norway rat (*Rattus norvegicus*), house mouse (*Mus musculus*), meadow vole (*M. pennsylvanicus*), muskrat (*Ondatra zibethicus*), meadow jumping mouse (*Zapus hudsonicus sic*), domestic dog (*Canis*), and domestic cat (*Felis*). Grimes and Beane (1984) assessed damage caused by beavers. Thompson (1994) caught house mice, western harvest mice, meadow voles, and fox squirrels in 240 trapnights (house mice represented 83% of the individuals captured), and also saw cottontails, muskrats, and beavers.

The purpose of our study was to document the presence and habitat use of birds and mammals in the first year-round, systematic survey of the South Platte urban corridor. We assessed how avian and mammalian species composition differed among different plant communities and compared the faunas with those re-

ported from other urban sites in the Denver area. We selected six permanent study areas in order to establish benchmarks for long-term study, to facilitate repeatable surveys in the future, and to allow establish-

ment of “living laboratories” where citizens can learn about riparian ecology. Results will allow optimization of human use of the river while minimizing negative impacts on wildlife.

MATERIALS AND METHODS

Major plant communities along the Platte consist primarily of cottonwood (*Populus deltoides*), elms (*Ulmus americanus* and *U. pumilus*), native grasses (including areas reseeded to native species), exotic grasses (mostly Kentucky bluegrass, *Poa pratensis*), weeds, and “other,” all in patches of various sizes (Instream Issues Task Force, 1996). General predictions can be made regarding the importance of some of these communities to wildlife (e.g., Bottorff, 1974). We chose not to sample the fauna in proportion to the representation of the various plant communities, which are being changed through restoration and other activities. It was difficult to find sites sufficiently large and continuous for sampling, due to the fragmentation of both native and non-native vegetation resulting from channelization, bank stabilization with riprap, and other extensive alterations. We chose six sites (Figure 1) so as to have two transects per site, with each site representing one of six community types (mesic, restored mesic, riparian woodland, non-native grassland, semi-native grassland, and urban pond):

Mesic community type, Riverside Cemetery: Our study site in mesic lowland is located downhill from a 145-acre cemetery that was established in 1876 (Kirkpatrick, 1998); the opposite bank is occupied by industry. The study site, partially coinciding with that used by Thompson (1994), is dominated by sandbar willow (*Salix exigua*), smooth brome (*Bromus inermis*), and sand dropseed (*Sporobolus cryptandrus*). Various trees include hawthorn (*Crataegus erythropoda*), Siberian elm, and cottonwoods. Most of the herbaceous vegetation is the result of a wetland restoration project conducted in 1993 (Thompson, 1994).

Restored mesic community type, Elitch Gardens: This site is an urban park adjacent to Six Flags Elitch Gardens Amusement Park, upstream of where Auraria was founded in 1858 (Jones and Forrest, 1973;




Figure 1. The six study sites along the urban South Platte River in Denver County, Colorado. Map by Adrian Kropp.

Kirkpatrick, 1998). The community includes sandbar willow, mustards (*Brassica* sp.), and dandelions (*Taraxacum officinale*), dominated by smooth brome (*B. inermis*) and other grasses. Saplings (*Populus* and

Ulmus) were planted at this site; there are no large trees, unlike the mesic community at Riverside Cemetery.

Riparian woodland community type, Grant-Frontier Park: An urban park (primarily planted to *P. pratensis*), a residential area, and industrial sites surround this small woodland. Most trees represent Siberian elm, with other conspicuous species including American elm, box elder (*Acer negundo*), wild rose (*Rosa woodsii*), and snowberry (*Symphoricarpos occidentalis*). The site is steeply sloped and thickly covered with grass, primarily *Bromus*; fallen logs, branches, and human litter also provide groundcover.

Non-native grassland community type, Johnson Habitat Park: This six-acre park, surrounded by industry, was formerly a city dump "slated to become an outdoor classroom amidst a number of restored natural environments" (Shoemaker, 1981:5). We conducted our research on an area including native blue grama (*Bouteloua gracilis*) but dominated by non-native cheatgrass, crested wheatgrass (*Agropyron cristatum*), diffuse knapweed (*Centaurea diffusa*), and white sweetclover (*Melilotus alba*). This field includes five small junipers (*Juniperus* cf. *scopulorum*) and virtually no vertical structure other than these trees; the field was mowed at least twice during our study.

Semi-native grassland community type, Overland Pond: This site is surrounded by an urban park and a golf driving range (Shoemaker 1981; Kirkpatrick, 1998). Two ponds were developed from an old gravel quarry, and the surrounding area was replanted with indigenous species such as rabbitbrush (*Chrysothamnus nauseosus*) in 1986. The plant community includes cheatgrass, blue grama, smooth brome, mullein (*Verbascum thapsus*), goldenpea (*Thermopsis rhombifolia*), sandbar willow, Siberian elm, and blue spruce (*Picea pungens*). Transects were located in grassland above the edge of the steep river escarpment, above the upper edge of the riparian woodland.

Urban pond, Globe Pond: Globe Pond is located near the northern county limit of Denver in the community of Globeville. Dominant vegetation includes cottonwoods (*Populus*), Siberian elm, cattails (*Typha*), cheatgrass (*Bromus tectorum*), and weedy forbs such as *Kochia scoparia*, *Cirsium*, and pigweed (*Amaranthus* spp.). This was the most insular of our sites, with

construction occurring outside of two of its boundaries during our study.

We used a combination of live-traps, scent stations, and transects to document the presence of as many birds and mammals as possible. We conducted sweep surveys of birds on each site following recommendations by Dawson (1981) and Ryder (1986). Permanent 300-meter line transects were established and marked with a permanent stake and geo-referenced using a global positioning system (GPS). All observers were experienced in identifying birds by both sight and sound. One person (Beane) walked the transect counting and identifying all birds observed or heard within 50 meters of the line. Two other observers conducted a sweep transect over each study area and counted all birds observed. Data were standardized by recording number of birds observed per linear transect and the amount of time per unit area. Surveys were conducted bimonthly over a 12-month period in 1998-99 (August, October, December, February, April, and June). Counts were conducted in the first five hours following sunrise. Each survey consisted of two counts on two consecutive days, with the order of the survey reversed on the second day.

We live-trapped small mammals using 7.5 x 7.5 x 22.5 cm (3"x3"x9") Sherman traps at 10 pace intervals, set in two transects per site with 15 traps per transect. We opened traps and baited them with oatmeal and peanut butter at or near sunset and checked traps at sunrise. We used Monel ear tags on rats and marked mice by toe-clipping (Nietfeld et al., 1996). We recorded identification, sex, and weight, with additional comments regarding reproductive condition. Traps were run for three consecutive nights in each of seven months (July, September, and November 1998, and January, March, May, and July 1999). Trapping effort was measured in trapnights (i.e., as the number of traps opened per night). Two Hav-a-hart traps also were employed at selected sites where the cover of vegetation sufficed to camouflage the traps.

One scent station was set up on open sand at each site to detect the presence of larger mammals. Each station consisted of a 1m² area of clear sand with predator scent left as bait in the middle of the sand. Stations were set up in the morning and checked 24 hours later.

Results.—Scientific names of vertebrates we identified are listed in Appendix A. We documented 65 species of birds (Table 1), including both migrants (such as the horned grebe, lesser yellowlegs, solitary sandpiper, western sandpiper) and winter residents (common goldeneye, herring gull, ring-billed gull, tree sparrow). The hooded merganser breeds in Colorado (Kingery, 1999) but was a winter resident on the South Platte. Of the 65 species, nine accounted for nearly 20% or more of the observations during each sampling season (Table 2).

Trapping effort totaled 1,890 trapnights consisting of 315 trapnights per site. The use of Sherman

traps documented the presence of five species of rodents (*R. megalotis*, *P. maniculatus*, *M. pennsylvanicus*, *R. norvegicus*, and *M. musculus*). Animal sightings, scat, and tracks at scent stations revealed the presence of ten additional species (Table 3). Magpies (*P. pica*) were the only non-target captures in Sherman and Hav-a-hart traps. There were very few recaptures. Three incidental mortalities (rodents) were prepared as vouchers and deposited in the Denver Museum of Nature and Science.

Discussion.—The elimination of native short-grass steppe and the establishment of various plant communities presumably changed the composition of

Table 1. Birds documented at Elitch's (E), Globe Pond (G), Grant Frontier (GR), Johnson Habitat Park (J), Overland Pond (O), and Riverside (R). Non-native species are marked with an asterisk.

	E (restored mesic)	G (urban pond)	GR (riparian)	J (disturbed grassland)	O (semi-native grassland)	R (mesic)
Horned grebe		x				
Pied-billed grebe		x	x			x
American white pelican		x				x
Double-crested cormorant	x	x	x	x	x	x
Great blue heron	x	x				x
Black-crowned night-heron		x	x	x		x
Green-backed heron						x
Domestic goose			x			
Canada goose	x	x	x	x	x	x
Gadwall		x	x	x	x	x
American wigeon		x	x		x	x
Mallard	x	x	x	x	x	x
Blue-winged teal			x		x	
Northern shoveler		x	x	x	x	x
Green-winged teal		x	x	x	x	x
Common goldeneye			x	x		
Hooded merganser			x			
Cooper's hawk					x	
Swainson's hawk			x			
Red-tailed hawk						x
American kestrel	x			x		x
American coot		x	x	x	x	
Killdeer	x	x		x	x	x
American avocet		x				x
Common snipe					x	
Lesser yellowlegs						x
Solitary sandpiper		x				
Western sandpiper		x				
Herring gull	x		x	x	x	
Ring-billed gull	x	x	x	x	x	x
Rock dove	x	x	x	x	x	x
Mourning dove		x	x	x		x
Great horned owl		x				
Belted kingfisher	x	x	x	x	x	x
Downy woodpecker			x	x		x

Table 1. (cont.)

	E (restored mesic)	G (urban pond)	GR (riparian)	J (disturbed grassland)	O (semi-native grassland)	R (mesic)
Northern flicker	x	x	x		x	x
Western wood-pewee		x	x			
Say's phoebe			x			
Loggerhead shrike						x
Plumbeous vireo			x			
Black-billed magpie	x	x	x	x	x	
American crow	x		x	x	x	x
Common raven	x		x	x	x	
Cliff swallow	x	x	x	x		x
Barn swallow	x	x	x	x	x	x
Black-capped chickadee		x	x	x	x	x
House wren					x	
American robin	x	x	x	x	x	x
Gray catbird		x				
European starling*	x	x	x	x	x	x
Yellow warbler		x	x		x	
Yellow-rumped warbler	x					
Lark sparrow		x		x		x
Song sparrow	x	x	x		x	x
White-crowned sparrow	x	x				x
Dark-eyed junco	x	x	x		x	
Red-winged blackbird	x	x	x	x	x	x
Brewer's blackbird			x		x	x
Common grackle	x	x	x	x	x	x
Bullock's oriole						x
House finch	x	x	x	x	x	x
American goldfinch		x				
American tree sparrow		x				x
House Sparrow*	x	x	x	x	x	x
Number of species/site:	23	46	37	32	34	41

the fauna along the South Platte. In rural Weld County, Colorado, Fitzgerald (1978) documented 109 species of birds and 24 species of mammals along the South Platte River, including prairie falcons (*Falco mexicanus*), horned larks (*Eremophila alpestris*), Ord's kangaroo rats (*Dipodomys ordii*), spotted ground squirrels (*Spermophilus spilosoma*), and other taxa associated with shortgrass steppe. We did not find these species or such large numbers of species. We found the fewest mammals on the non-native grassland and the fewest birds on the restored mesic site, both of which lack mature elms and cottonwoods (Tables 1, 3). The greatest number of species occurred on two sites with vegetation providing well-developed vertical structure (Globe Pond and the lowland mesic site at Riverside Cemetery).

We saw birds primarily on the river or in trees along riverbanks, and the species listed are typical of riparian communities. For example, birds such as the common goldeneye are typical of open water, whereas the green heron prefers riparian woods (Andrews and Righter, 1992). This region of the South Platte supports large numbers of ducks in winter; for example, for the 1999-2000 Christmas Bird Count period, 16 species of ducks were recorded, comprising 36.4% of the count results (H. E. Kingery, pers. comm., 2000). We detected all of the 13 avian species known to breed in lowland riparian ecosystems in Colorado (Preston and Kingery, 1998), except for western and eastern kingbirds (*Tyrannus verticalis* and *T. tyrannus*) and blue grosbeak (*Guiraca caerulea*). Of the 13 species found at all 6 sites (Table 1), the mallard, double-

Table 2. Dominant bird species observed by season and location.

	Grant Frontier	Overland Pond	Johnson Habitat	Elitches	Globe Pond	Riverside
Late Summer - August % of Observations	None	Canada Goose 39.0%	House Finch 28.8%	None	Canada Goose 23%	None
Fall - October % of Observations	Mallard 40.7%	None	Green-winged Teal 45.6%	Canada Goose 42.1%	House Finch 36%	House Finch 29.1%
Winter - December % of Observations	Canada Goose 48.1%	Canada Goose 66.7%	Ring-billed Gull 48.6%	Ring-billed Gull 76.5%	Ring-billed Gull 32.8%	European Starling 45.2%
Winter - February % of Observations	Mallard 23.8%	Gadwall 19.4%	Ring-billed Gull 26.2%	Common Grackle 33.9%	European Starling 19.6%	Canada Goose 30.3%
Spring - April % of Observations	Gadwall 30.5%	Gadwall 26.4%	Gadwall 29.7%	Ring-billed Gull 32.7%	Common Grackle 29.7%	White Pelican 29.9%
Summer - June % of Observations	European Starling 26.3%	Common Grackle 32.4%	Canada Goose 38.6%	Common Grackle 18.2%	European Starling 26.3%	Common Grackle 28.3%

Table 3. Mammals documented at sites as indicated on Table 1. Non-native species are marked with an asterisk. Abundance codes are from the Colorado Natural Diversity Information Source (2000).

Site	E (restored mesic)	G (urban pond)	GR (riparian)	J (disturbed grassland)	O (semi-native grassland)	R (lowland mesic)	Abundance
Cottontail		x		x	x		fairly common
Eastern fox squirrel	x			x			common
American beaver			x			x	fairly common
Western harvest mouse		x				x	fairly common
Deer mouse		x				x	abundant
Meadow vole					x		common
Common muskrat		x					common
Norway rat *	x		x			x	common
House mouse*	x					x	abundant
Dog*	x		x				common
Coyote		x					common
Red fox	x				x		abundant
Common raccoon	x		x		x		abundant
Cat*							fairly common
Deer				x		x	fairly common
Number of species detected:	7	8	7	4	8	9	
Number of captures of rodents:	61	40	16	0	17	72	

crested cormorant, belted kingfisher, and red-winged blackbird also indicate the presence of water. The remaining ten (Canada goose, ring-billed gull, rock dove, barn swallow, American robin, European starling, common grackle, house finch, and house sparrow) are widespread birds common to urban areas (e.g., Loeffler, 1990, Kingery, 1998b; Colorado NDIS, 2000). The rock dove, European starling, and house sparrow are exotics introduced into the United States in the 18th and 19th centuries; they now comprise up to 95% of some urban bird communities during winter (Adams, 1994).

The dominant birds also were primarily urban-adapted species (Table 2). Six of the nine dominant species (Canada goose, mallard, ring-billed gull, European starling, common grackle, and house finch) are common urban birds, whereas the remaining three are migrant waterfowl (green-winged teal, gadwall, and white pelican). Most observations of these waterfowl were recorded in spring, late winter, or fall. Canada geese and ring-billed gulls were the most common wintering species at all sites, whereas the most common summer species at four of the six sites was the common grackle (Table 2).

Raptors are important indicators of ecosystem health and especially popular among bird-watchers. A great horned owl, Cooper's hawks, Swainson's hawks, and red-tailed hawks all were present, and all are uncommon sights in Denver (Colorado NDIS, 2000). The American kestrel was the most widely-reported raptor (Table 1). Its apparent success along the South Platte River might be attributed to its small size (it maintains the smallest territories of all Colorado hawks) and versatile diet (Winn, 1998).

The mammalian community was similar to that reported for other local urban areas. Two cottontails (*S. audubonii* and *S. floridanus*) are fairly common, but difficult to identify without a specimen in hand (Fitzgerald et al., 1994; Colorado NDIS, 2000). We detected deer sign at only one site (Table 3); whitetails (*O. virginianus*) are more closely associated with riparian regions than mule deer (*O. hemionus*), but both species are fairly common (Colorado NDIS, 2000). Of the native rodents encountered, the American beaver and western harvest mouse are considered fairly common, the deer mouse abundant, and the meadow

vole and muskrat common in Denver (Colorado NDIS, 2000). Western harvest mice occur in wetlands, weedy or grassy areas, and riparian woodlands throughout eastern Colorado (Fitzgerald et al., 1994). The deer mouse is the most ubiquitous of North American rodents, using a wide variety of habitats and adapting particularly well to disturbance (e.g., Fitzgerald et al., 1994; Handley, 1999). Meadow voles prefer mesic areas but tolerate patchy habitats and edges well (e.g., Harper et al., 1993; Fitzgerald et al., 1994). Beaver and muskrat occur throughout Colorado where there are permanent bodies of water (Fitzgerald et al., 1994).

We expected to find the three widespread, non-native rodents observed (Table 3). Fox squirrels were introduced in several places in Colorado and have become common watchable wildlife in Denver and many other cities (e.g., Loeffler, 1990; Adams, 1994; Flyger, 1999). The Old World Norway rat and house mouse are cosmopolitan. The only site at which we did not capture *Mus* was the disturbed grassland, which lacked significant ground cover.

Raccoons were documented at four locations, but probably occur at all sites. They are very common in both urban and riparian areas and appear to be increasing in numbers in this region (Fitzgerald et al., 1994; VanDruff et al., 1996). The most common wild carnivore was the red fox, the most widely distributed wild carnivore in the world (Seidensticker, 1999). We detected the coyote at only one site, but it is common in Denver (Table 3). We saw domestic dogs at all six sites, and at least some of these dogs were living in dens and appeared to be feral. We found evidence of house cats at two locations. Domestic dogs and cats both have been implicated in the decline and disappearance of native species, although their impacts on wildlife have been poorly studied (Atkinson, 1989; Adams, 1994; Clarke and Pacin, 2002).

No animals were captured in transects on the disturbed, non-native grassland, which reflected the lack of vertical structure due to frequent mowing. This site also had the lowest species richness. The greatest numbers of rodent captures occurred in the two mesic areas (Table 3). Harvest mice were the most frequently-captured rodents in true mesic lowland. *Mus musculus* was the only mouse caught in the restored mesic community until 9 July 1999, when three meadow voles

were trapped. The presence of voles might indicate community succession following restoration efforts less than three years old, or might indicate a change in interspecific competition; interactions between *Mus* and native species should be examined.

Meadow voles, house mice, deer mice, Norway rats, and western harvest mice have been reported from elsewhere in the metropolitan area (Jones, 1997; ERO Resources, 1998). The presence of additional species at other nearby urban sites suggested that the South Platte mammalian fauna was somewhat depauperate. Both the prairie vole (*M. ochrogaster*) and the Mexican wood rat (*Neotoma mexicana*) have been reported from elsewhere in Denver County (Beane and Powell, 1997; Jones, 1997).

The absence of *Zapus* was of special note. Its presence along the urban South Platte was documented by a capture in "waist-high forbs over moist ground" (Erickson, 1982) at a site that is now the location of a children's museum and urban park lacking the original multi-stratum plant community. The only subspecies of jumping mouse (*Z. hudsonius preblei*) occurring in the area was listed as federally threatened (U.S. Fish and Wildlife Service, 1998). Most records are from moist riparian areas with thick herbaceous vegetation (Jones, 1999, and citations therein). Its absence along the urban South Platte indicated the lack of extensive native vegetation along this stretch of river.

Urban river corridors are popular locales for recreation, including watching wildlife. In Colorado, outdoor activities are considered tremendously important, as evidenced by recent surveys regarding preferences regarding wildlife viewing (Manfredo et al., 1991) and use of nature trails (Cuciti, 2002), and by large numbers of volunteers assisting in projects such as the breeding bird atlas (Kingery, 1998a) and bat surveys (Navo et al., 2001). In the Denver metropolitan area, the history of the Rocky Mountain Arsenal and other sites demonstrates the high level of public interest in natural history, although local governments strongly support economic growth and development (Jones and Preston, 2000). The lowland riparian ecosystem offers unique opportunities for public education, in spite of representing only about 3% of Colorado's area. This ecosystem supports a relatively high number of mammalian species, more native birds than the shortgrass

steppe, and is one of the most threatened of western ecosystems (Cooperrider, 1989; Fitzgerald et al., 1994; Ohmart, 1994; Preston and Kingery, 1998; Fleischner, 1999). Additionally, riparian corridors are thought to serve as avenues of dispersal, particularly for species from the east, although the historic influence on dispersal remains unknown (e.g., Fitzgerald et al., 1994; Kingery, 1998b). Other positive values connected with the presence of urban plants and animals include aesthetic, environmental (e.g., services such as clean water and noise abatement), economic, psychological, and sociological benefits (Adams and Dove, 1989; Loeffler, 1990; Moorhead, 1992; Schwarz, 1993; Botkin and Beveridge, 1997; Werner, 1999). We make the following recommendations for management of urban areas, particularly those in riparian zones:

Maintenance and Management

The appearance of most urban recreational areas is maintained in a manner considered appropriate by most citizens. For example, in 1999, efforts to maintain the aesthetic appearance of the South Platte corridor included the equivalent of 9.5 miles of tree trimming and 199 miles of trash and debris pickup and removal (Urbonas, 1999). However, litter in the form of fallen trees and branches should be left for species known to respond positively to vertical structure (Douglass and Reinert, 1982; Getz, 1985; Sears and Anderson, 1991; Adams, 1994; VanDruff et al., 1996). Stable rodent populations also are key determinants in the distribution of very popular raptors (e.g., Preston and Beane, 1996). Considerations for public safety must be balanced with the needs of wildlife. Education might help modify current aesthetic values.

Special efforts should be made to manage for wildlife on less disturbed areas than on more frequently visited sites. Traffic noise negatively affects densities of breeding birds (Reijnen et al., 1997). Starlings and other less-desired generalists adapt better to human disturbance than do specialists (Hallock, 1989; Sears and Anderson, 1991; VanDruff et al., 1996). Many species are impacted directly and indirectly by roads (Trombulak and Frissell, 2000).

The restoration of native vegetation should continue for both aesthetic and biological reasons. Botkin and Beveridge (1997) offered recommendations for

managing for the best mix of urban vegetation types related to water use, and it has been reported that native birds in the Rocky Mountain region use native trees more than non-native birds (Sears and Anderson, 1991).

Put in place a mechanism, such as a long-term management plan, to ensure that long-term restoration efforts survive changes in politics and personnel. For example, on one of our sites, the restoration of xeric plants has not resulted in establishment of a xeric ecosystem because of how the park was managed subsequently.

Education

Institutions managing parks should mandate ongoing educational efforts, including on-site signage and addresses of pertinent websites such as those of Ballenger (1996a, 1996b), to ensure that the public has ready access to up-to-date information.

The widespread distribution of alien species requires that these organisms be of special importance in education and research. Herein we have treated non-natives as 'exotics'. However, Haspel and Calhoon (1991:27) stated that the study of urban cats "...is the study of a species in a unique ecosystem, one in which it has been reproductively successful for thousands of years." Such intriguing ideas should be investigated for development of research and educational programming. How do invasive species interact with native wildlife in urban areas? In any case, exotics need to be included in any educational programming.

Projects such as the development of the South Platte Greenway provide opportunities to provide participation by local residents. Including residents in planning and incorporating information about local culture and history will promote alliances between city planners, environmentalists, and advocates of social justice (Chapman, 2001).

Research

Encourage additional research and peer-reviewed publications regarding urban areas. Many valuable data lie buried in 'gray literature', as in the report by Thompson (1994) and other reports cited in this paper. Various projects have been conducted regarding envi-

ronmental quality (for example, see Kimbrough and Litke, 1996; Saldiva and Böhm, 1998) but more ecological work needs to be done regarding the ecology unique to cities. People in urban ecosystems can weaken or enhance the natural forces affecting ecosystems (Pickett and McDonnell, 1993; Collins et al., 2000). Such human-induced effects in Colorado include the expansion of house sparrow populations to elevations exceeding 3,000 m or 10,000 ft (Kaempfer, 1998). Some popular articles have treated ecological subjects (e.g., Kerlinger, 1999; Jones, 2000), and various examples of technical publications regarding urban ecology include those by Hallock (1989), Hiebert (1990), Groom (1993), Lajeunesse et al. (1997), Reijnen et al. (1997), Fernández-Juricic (2000), and Grindler and Krausman (2001). However, results of a recent survey demonstrated that very few papers in ecological journals dealt with urban biology (Collins et al., 2000).

The relative paucity of ecological research in urban areas counters widespread needs for biological input in urban planning (Murphy, 1988; Botkin and Beveridge, 1997; Babbitt, 1999) and is being partially addressed through establishment of the US Long-Term Ecological Research network (Parlange, 1998; Grimm et al., 2000). The South Platte is one of those urban sites providing a unique opportunity in the study of biological events that differ, or occur on a different time scale, than those in less impacted areas. We know that dramatic changes in floras and faunas have occurred since European settlement. In most cities, we lack data regarding the flora and fauna present before urban development. However, surveys such as ours provide baseline data and the opportunity to document alterations in bird and mammal communities as future changes occur along urban river corridors and other urban sites.

ACKNOWLEDGMENTS

We thank volunteers L. M. Carpenter, J. Harting, and C. Stucheli, and an anonymous reviewer. A. Kropp provided the map, R. Brune shared unpublished vegetation data, H. E. Kingery assisted with bird records, and B. Chadwick served as our liaison with the city. This project was supported by the Mayor's South Platte River Commission and by the Denver Museum of Nature and Science.

LITERATURE CITED

- Adams, L. W. 1994. Urban wildlife habitats: a landscape perspective. *Wildlife Habitats Volume 3*, University of Minnesota Press, Minneapolis, 186 pp.
- Adams, L. W., and L. E. Dove. 1989. Wildlife reserves and corridors in the urban environment: a guide to ecological landscape planning and resource conservation. National Institute for Urban Wildlife, Shepherdstown, West Virginia, 91 pp.
- Alston, M. 1999. Denver's South Platte River Corridor: projects progress towards a vision, lots left to do. U.S. Environmental Protection Agency, Community Based Environmental Protection Summer 1999 News, www.epa.gov/region08.
- Andrews, R., and R. Righter. 1992. Colorado birds: a reference to their distribution and habitat. Denver Museum of Natural History, Denver, Colorado, 442 pp.
- Anonymous. 1997. International chorus day Sunday, May 4, 1997. Colorado Urban Wildlife Partnership Newsletter.
- Anonymous. 1998. Greenway vision saves river and generates investment. Urban Parks Institute, www.pps.org/urbanparks.
- Atkinson, I. 1989. Introduced animals and extinctions. Pages 54-75, in *Conservation for the twenty-first century* (D. Western and M. C. Pearl, eds.), Wildlife Conservation International, New York, New York, 365 pp.
- Babbitt, B. 1999. Noah's mandate and the birth of urban bioplanning. *Conservation Biology*, 13:677-678.
- Ballenger, L. 1996a. *Mus musculus*. www.ummz.umich.edu.
- Ballenger, L. 1996b. *Rattus norvegicus*. www.ummz.umich.edu.
- Beane, R. D., and M. Powell. 1997. Preble's meadow jumping mouse presence/absence surveys, Bluff Lake Natural Area, Denver County, Colorado. Unpublished report prepared for Stapleton Development Corporation, Denver, Colorado.
- Botkin, D. B., and C. E. Beveridge. 1997. Cities as environments. *Urban Ecosystems*, 1:3-19.
- Bottorff, R. L. 1974. Cottonwood habitat for birds in Colorado. *American Birds*, 28:975-979.
- Brenneman, B. 1973. Miracle on Cherry Creek: an informal history of the birth and re-birth of a neighborhood. World Press Inc., Denver, Colorado, 130 pp.
- Brittan, M. 1992. South Platte Park small mammal research. Unpublished report, South Platte Park, 6 pp.
- Brovksy, C. 1999. New life comes to central Platte. *The Denver Post* August 1, B1, B7.
- Chapman, G. 2001. The intersection of environmental planning and social justice: Denver's Platte River Greenway. Master in City Planning Thesis, Massachusetts Institute of Technology, Cambridge, Massachusetts.
- City and County of Denver, Office of the Mayor. 2000. South Platte River urban watershed restoration program. Briefing document.
- Clarke, A. L., and T. Pacin. 2002. Domestic cat "colonies" in natural areas: a growing exotic species threat. *Natural Areas Journal*, 22:154-159.
- Collins, J. P., A. Kinzing, N. B. Grimm, W. F. Fagan, D. Hope, J. Wu, and E. T. Boner. 2000. A new urban ecology. *American Scientist*, 88:416-425.
- Colorado Natural Diversity Information Source. 2000. www.nds.nrel.colostate.edu.
- Cooper, A., A. Armstrong, and C. Kampert. 1990. The WildWatch book. Denver Museum of Natural History in cooperation with Roberts Rinehart Publishers, Niwot, Colorado, 90 pp.
- Cooperrider, A. 1989. Conservation of biological diversity in the west. Pages 3-9 in *Issues and technology on the management of impacted wildlife: proceedings of a national symposium*, Glenwood Springs, Colorado. (P. R. Davis, J. C. Emerick, D. M. Finch, S. Q. Foster, J. W. Monarch, S. Rush, O. Thorne II, and J. Todd, eds), Thorne Ecological Institute, Boulder, Colorado, 213 pp.
- Cuciti, P. 2002. Results of Colorado statewide public survey on attitudes toward trails. www.americantrails.org.
- Dawson, D. G. 1981. Experimental design when counting birds. Pages 392-398 in *Studies in avian biology* no. 6, (C. J. Ralph and M. S. Scott, eds.), Cooper Ornithological Society, 630 pp.
- Douglass, N. J., and H. K. Reinert. 1982. The utilization of fallen logs as runways by small mammals. *Proceedings of the Pennsylvania Academy of Science*, 56:162-164.
- Erickson, D. L. 1982. Survey of plant and mammal life along the Denver Platte River Greenway. Unpublished report, funded by the Denver Audubon Society.

- ERO Resources Corporation. 1998. Preble's meadow jumping mouse trapping survey results for Cherry Creek at Monaco Street, Denver County, Colorado. Unpublished report prepared for Nolte and Associates, by ERO Resources Corporation, 1842 Clarkson Street, Denver, Colorado.
- Fernández-Juricic, E. 2000. Avifaunal use of wooded streets in an urban landscape. *Conservation Biology*, 14:513-521.
- Fitzgerald, J. P. 1978. Vertebrate associations in plant communities along the South Platte River in north-eastern Colorado. Pages 73-88 in *Lowland river and stream habitat in Colorado: a symposium* (W. D. Graul and S. J. Bissell, technical coordinators), Colorado Chapter of the Wildlife Society and Colorado Audubon Council, Greeley, Colorado, 195 pp.
- Fitzgerald, J. P., C. A. Meaney, and D. M. Armstrong. 1994. *Mammals of Colorado*. Denver Museum of Natural History and University Press of Colorado, Niwot, Colorado, 467 pp.
- Fleischner, T. L. 1999. Keeping the cows off: conserving riparian areas in the American west. Pages 64-65, in *Terrestrial ecosystems of North America: a conservation assessment* (T. H. Ricketts, E. Dinerstein, D. M. Olson, C. J. Loucks, W. Eichbaum, D. DellaSala, K. Kavanagh, P. Hedao, P. T. Hurley, K. M. Karney, R. Abell, and S. Walters, eds.), Island Press, Washington, D. C., 485 pp.
- Flyger, V. 1999. Eastern fox squirrel *Sciurus niger*. Pages 456-458 in *The Smithsonian book of North American mammals* (D. E. Wilson and S. Ruff, eds.), Smithsonian Institution Press, Washington, D.C., 750 pp.
- Getz, L. L. 1985. Habitats. Pages 286-309 in *Biology of New World Microtus* (R. H. Tamarin, ed.), American Society of Mammalogists Special Publication No. 8, 893 pp.
- Grimes, M. D., and R. D. Beane. 1984. The effect of beaver on the South Platte River. Unpublished report submitted to the Platte River Greenway Foundation, 1666 S. University Blvd., Denver, Colorado, 15 pp.
- Grimm, N. B., J. M. Grove, S. T. A. Pickett, and C. L. Redman. 2000. Integrated approaches to long-term studies of urban ecological systems. *BioScience*, 50:571-584.
- Grinder, M. I., and P. R. Krausman. 2001. Home range, habitat use, and nocturnal activity of coyotes in an urban environment. *Journal of Wildlife Management*, 65:887-898.
- Groom, D. W. 1993. Magpie *Pica pica* predation on black-bird *Turdus merula* nests in urban areas. *Bird Study*, 40:55-62.
- Hallock, D. 1989. Wildlife trends on the urban-rural fringe. Pages 17-22, in *Issues and technology on the management of impacted wildlife: proceedings of a national symposium*, Glenwood Springs, Colorado. (P. R. Davis, J. C. Emerick, D. M. Finch, S. Q. Foster, J. W. Monarch, S. Rush, O. Thorne II, and J. Todd, eds.), Thorne Ecological Institute, Boulder, Colorado, 213 pp.
- Handley, C. O., Jr. 1999. Deer mouse *Peromyscus maniculatus*. Pages 575-577, in *The Smithsonian book of North American mammals* (D. E. Wilson and S. Ruff, eds.), Smithsonian Institution Press, Washington, D.C., 750 pp.
- Harper, S. J., E. K. Bollinger, and G. W. Barrett. 1993. Effects of habitat patch shape on population dynamics of meadow voles (*Microtus pennsylvanicus*). *Journal of Mammalogy*, 74:1045-1055.
- Haspel, C., and R. E. Calhoon. 1991. Ecology and behavior of free-ranging cats in Brooklyn, New York. Pages 27-30 in *Wildlife conservation in metropolitan environments* (L. W. Adams and D. L. Leedy, eds.), National Institute for Urban Wildlife, Columbia, Maryland, 264 pp.
- Hekkers, J., editor. 1987. *Wild in the streets: Denver area wildlife adjusts to urban living*. Colorado Division of Wildlife, Publication DOW-M-1-2-87, Denver, Colorado, 19 pp.
- Hiebert, R. D. 1990. An ecological restoration model: application to razed residential sites. *Natural Areas Journal*, 10:181-186.
- Instream Issues Task Force. 1996. *South Platte River Corridor Project: instream issues report*, 92 pp.
- Johnson, W. C. 1994. Woodland expansions in the Platte River, Nebraska: patterns and causes. *Ecological Monographs*, 64:45-84.
- Jones, A. L. 1997. Presence/absence surveys for Preble's meadow jumping mouse: Sand Creek---Havana and Peoria crossings. Unpublished report prepared for Stapleton Development Corporation by MDG, Inc., 820 Santa Fe Drive, Denver, Colorado.
- Jones, C. A. 1999. *Zapus hudsonius* in southern Colorado. *Occasional Papers, Museum of Texas Tech University*, 191:1-7.
- Jones, C. A. 2000. A river runs through it. *Museum Quarterly, Denver Museum of Nature and Science*, 9(2): 2-5.

- Jones, C. A., and C. R. Preston. 2000. The role of the bald eagle as a flagship species for the Rocky Mountain Arsenal: from Superfund Site to wildlife refuge. Pages 165-171 in *Reflections of a naturalist: papers honoring Professor Eugene D. Fleharty*. (J. R. Choate, ed.), Fort Hays Studies, Fort Hays State University, Special Issue 1, 241 pp.
- Jones, W. C., and K. Forrest. 1973. Denver: a pictorial history from frontier camp to queen city of the plains. Pruett Publishing Company, Boulder, Colorado, 336 pp.
- Kaempfer, B. 1998. House sparrow *Passer domesticus*. Pages 542-543 in *Colorado breeding bird atlas* (H. E. Kingery, ed.), Colorado Bird Atlas Partnership and Colorado Division of Wildlife, Denver, Colorado, 636 pp.
- Kerlinger, P. 1999. Nesting in Gotham. *Natural History*, June 1999: 24-26.
- Kimbrough, R. A., and D. W. Litke. 1996. Pesticides in streams draining agricultural and urban areas in Colorado. *Environmental Science and Technology*, 30:908-916.
- Kingery, H. E., editor. 1998a. *Colorado breeding bird atlas*. Colorado Bird Atlas Partnership and Colorado Division of Wildlife, Denver, Colorado, 636 pp.
- Kingery, H. E. 1998b. Post-settlement changes to Colorado habitats. Pages 33-35 in *Colorado breeding bird atlas* (H. E. Kingery, ed.), Colorado Bird Atlas Partnership and Colorado Division of Wildlife, Denver, Colorado, 636 pp.
- Kingery, H. E. 1999. Hooded merganser breeding in Colorado. *Journal of the Colorado Field Ornithologists*. 33:180-182
- Kingery, H. E. (count leader). 2000. Denver urban Christmas bird count. *The Lark Bunting* 35(2):5-12.
- Kirkpatrick, T. 1998. *The South Platte River Greenway Trail: the trail user's guide*. Trails 'N Tales Publishing Co., Englewood, Colorado, 82 pp.
- Lajeunesse, D., G. Domon, A. Cogliastro, and A. Bouchard. 1997. Monitoring recreational use in urban natural areas. *Natural Areas Journal*, 17:366-379.
- Loeffler, C. 1990. Taking a look at urban wildlife. Colorado Division of Wildlife, Publication DOW-M-13-88, Colorado Springs, Colorado, 42 pp.
- Manfredo, M. J., A. Bright, and M. Stephenson. 1991. Public preferences for non-consumptive wildlife recreation in the Denver area. Final report, submitted to the Colorado Division of Wildlife, Denver, Colorado, 73 pp.
- Mayor's South Platte River Commission. 1995. *Great Outdoors Colorado Legacy Grant Concept Paper*, 12 pp.
- Meister, G. E. 1965. Trajectory of a tragedy: Denver area flood, June 16th-17th, 1965. Hotchkiss Inc., Denver, Colorado, 30 pp.
- Miller, J. 1996. Urbanization and bird communities in riparian areas. *Urban Wildlife Newsletter* 25:2-3.
- Mitchell, R. S. 1972. Small rodents of the flood plain of the South Platte River at the proposed Narrows Reservoir site. M. A. thesis, University of Northern Colorado, Greeley, Colorado, 50 pp.
- Moorhead, C. A. 1992. *Colorado's backyard wildlife: a natural history, ecology, and action guide to Front Range urban wildlife*. Roberts Rinehart Publishers, Niwot, Colorado, 133 pp.
- Muntz, G., and A. Wuth. 1983. *A path through time: a guide to the Platte River Greenway*. Jende-Hagan Book Corporation, Frederick, Colorado.
- Murphy, D. D. 1988. Challenges to biological diversity in urban areas. Pages 71-76 in *Biodiversity* (E. O. Wilson, ed.), National Academy Press, Washington, D. C., 519 pp.
- Navo, K. W., T. E. Ingersoll, L. R. Bonewell, and N. LaMantia-Olson. 2001. *Bats/Inactive Mines Project, 2001 mine evaluations summary report*. Submitted to Great Outdoors Colorado, 7 pp.
- Nietfeld, M. T., N. W. Barrett, and N. Silvy. 1996. Wildlife marking techniques. Pages 140-168 in *Research and management techniques for wildlife and habitats* (T. A. Bookhout, ed.), The Wildlife Society, Bethesda, Maryland, 740 pp.
- Noel, T. J. 1980. *Denver: Rocky Mountain gold*. Continental Heritage Press, Tulsa, Oklahoma, 256 pp.
- Ohmart, R. D. 1994. The effects of human-induced changes on the avifauna of western riparian habitats. *Studies in Avian Biology*, 15:273-285.
- Olson, T. E., and F. L. Knopf. 1988. Patterns of relative diversity within riparian small mammal communities, Platte River watershed, Colorado. Pages 379-386 in *Management of amphibians, reptiles, and small mammals in North America: proceedings of the symposium* (R. C. Szaro, K. E. Severson, and D. R. Patton, technical coordinators), U.S. Department of Agriculture Forest Service General Technical Report RM-166, 458 pp.
- Parlange, M. 1998. The city as ecosystem. *BioScience*, 48:581-585.

- Pickett, S. T. A., and M. J. McDonnell. 1993. Humans as components of ecosystems: a synthesis. Pages 310-316 in *Humans as components of ecosystems: the ecology of subtle human effects and populated areas* (M. J. McDonnell and S. T. A. Pickett, eds.), Springer-Verlag, New York, New York, 364 pp.
- Preston, C. R., and R. D. Beane. 1996. Occurrence and distribution of diurnal raptors in relation to human activity and other factors at Rocky Mountain Arsenal, Colorado. Pages 365-375 in *Raptors in human landscapes: adaptations to built and cultivated environments* (D. M. Bird, D. E. Varland, and J. J. Negro, eds.), Academic Press, San Diego, California, 396 pp.
- Preston, C. R., and H. E. Kingery. 1998. The Colorado environment. Pages 21-32 in *Colorado breeding bird atlas* (H. E. Kingery, ed.), Colorado Bird Atlas Partnership and Colorado Division of Wildlife, Denver, Colorado, 636 pp.
- Reijnen, R., R. Foppen, and G. Veenbaas. 1997. Disturbance by traffic of breeding birds: evaluation of the effect and considerations in planning and managing road corridors. *Biodiversity and Conservation*, 6:567-581.
- Ryder, R. A. 1986. Songbirds. Pages 291-312 in *Inventory and monitoring of wildlife habitat* (A. Y. Cooperrider, R. J. Boyd, and H. R. Stuart, eds.), U.S. Department of Interior, Bureau of Land Management Service Center, Denver, Colorado, 858 pp.
- Saldiva, P. H. N., and G. M. Böhm. 1998. Animal indicators of adverse effects associated with air pollution. *Ecosystem Health*, 4:230-235.
- Schwarz, L. LaB. 1993. *Greenways: a guide to planning, design, and development*. The Conservation Fund, Island Press, Washington, D.C., 351 pp.
- Sears, A. R., and S. H. Anderson. 1991. Correlations between birds and vegetation in Cheyenne, Wyoming. Pages 75-80 in *Wildlife conservation in metropolitan environments* (L. W. Adams and D. L. Leedy, eds.), National Institute for Urban Wildlife, Columbia, Maryland, 264 pp.
- Seidensticker, J. 1999. Red fox *Vulpes vulpes*. Pages 150-152 in *The Smithsonian book of North American mammals* (D. E. Wilson and S. Ruff, eds.), Smithsonian Institution Press, Washington, D.C., 750 pp.
- Shoemaker, J. 1981. *Returning the Platte to the people: a story of a unique committee, the Platte River Development Committee*. The Greenway Foundation and Tumbleweed Press, Westminster, Colorado, 118 pp.
- Shuster, E. 1996. *The Platte River Greenway: the forgotten Nile of North America*. American Trails Resources and Library, www.outdoorlink.com/amtrails.
- Thompson, R. W. 1994. 1993 wildlife monitoring report: Riverside and Windsor Gardens Wetlands Fairlake Project. Unpublished report prepared for Fairwind Associates by Western Resource Development, Denver, Colorado, 52 pp.
- Tromulak, S. C., and C. A. Frissell. 2000. Review of ecological effects of roads on terrestrial and aquatic communities. *Conservation Biology*, 14:18-30.
- United States Environmental Protection Agency. 1999. *The South Platte River in Colorado*. EPA 908-F-98-002, Region 8 8EPR-EP. www.epa.gov/region8.
- United States Fish and Wildlife Service. 1998. Final rule to list the Preble's meadow jumping mouse as a threatened species. *Federal Register* 63, 92:1-28.
- Urbonas, B. 1999. South Platte River program notes. *Flood Hazard News* 29(1), unpaginated. www.udfcd.org.
- VanDruff, L. W., E. G. Bolen, and G. J. San Julian. 1996. Management of urban wildlife. Pages 507-530 in *Research and management techniques for wildlife and habitats* (T. A. Bookhout, ed.), The Wildlife Society, Bethesda, Maryland, 740 pp.
- Walker, J. 1997. Helpers green up S. Platte parks. *The Denver Post* May 4, 2B.
- Werner, P. 1999. Why biotope mapping in populated areas? *Deinsea*, 5:9-26.
- Winn, R. 1998. American kestrel. Pages 126-127 in *Colorado breeding bird atlas* (H. E. Kingery, ed.), Colorado Bird Atlas Partnership and Colorado Division of Wildlife, Denver, Colorado, 636 pp.

APPENDIX A

Scientific and vernacular names of birds and mammals reported in this study. Non-native species are marked with an asterisk.

Birds

Horned grebe	<i>Podiceps auritus</i>	American crow	<i>Corvus brachyrhynchos</i>
Pied-billed grebe	<i>Podilymbus podiceps</i>	Common raven	<i>Corvus corax</i>
American white pelican	<i>Pelecanus erythrorhynchos</i>	Cliff swallow	<i>Petrochelidon pyrrhonota</i>
Double-crested cormorant	<i>Phalacrocorax auritus</i>	Barn swallow	<i>Hirundo rustica</i>
Great blue heron	<i>Ardea herodias</i>	Black-capped chickadee	<i>Poecile atricapillus</i>
Black-crowned night-heron	<i>Nycticorax nycticorax</i>	House wren	<i>Troglodytes aedon</i>
Green-backed heron	<i>Butorides striatus</i>	American robin	<i>Turdus migratorius</i>
Domestic goose	<i>Anser anser*</i>	Gray catbird	<i>Dumetella carolinensis</i>
Canada goose	<i>Branta canadensis</i>	European starling	<i>Sturnus vulgaris*</i>
Gadwall	<i>Anas strepera</i>	Yellow warbler	<i>Dendroica petechia</i>
American wigeon	<i>Anas americana</i>	Yellow-rumped warbler	<i>Dendroica coronata</i>
Mallard	<i>Anas platyrhynchos</i>	Lark sparrow	<i>Chondestes grammacus</i>
Blue-winged teal	<i>Anas discors</i>	Song sparrow	<i>Melospiza melodia</i>
Northern shoveler	<i>Anas clypeata</i>	White-crowned sparrow	<i>Zonotrichia leucophrys</i>
Green-winged teal	<i>Anas crecca</i>	Dark-eyed junco	<i>Junco hyemalis</i>
Common goldeneye	<i>Bucephala clangula</i>	Red-winged blackbird	<i>Agelaius phoeniceus</i>
Hooded merganser	<i>Lophodytes cucullatus</i>	Brewer's blackbird	<i>Euphagus cyanocephalus</i>
Cooper's hawk	<i>Accipiter cooperii</i>	Common grackle	<i>Quiscalus quiscula</i>
Swainson's hawk	<i>Buteo swainsoni</i>	Bullock's oriole	<i>Icterus bullockii</i>
Red-tailed hawk	<i>Buteo jamaicensis</i>	House finch	<i>Carpodacus mexicanus</i>
American kestrel	<i>Falco sparverius</i>	American goldfinch	<i>Carduelis tristis</i>
American coot	<i>Fulica americana</i>	American tree sparrow	<i>Spizella arborea</i>
Killdeer	<i>Charadrius vociferus</i>	House sparrow	<i>Passer domesticus*</i>
American avocet	<i>Recurvirostra americana</i>		
Common snipe	<i>Gallinago gallinago</i>		
Lesser yellowlegs	<i>Tringa flavipes</i>		
Solitary sandpiper	<i>Tringa solitaria</i>		
Western sandpiper	<i>Calidrus mauri</i>		
Herring gull	<i>Larus argentatus</i>		
Ring-billed gull	<i>Larus delawarensis</i>		
Rock dove	<i>Columba livia*</i>		
Mourning dove	<i>Zenaida macroura</i>		
Great horned owl	<i>Bubo virginianus</i>		
Belted kingfisher	<i>Ceryle alcyon</i>		
Downy woodpecker	<i>Picoides pubescens</i>		
Northern flicker	<i>Colaptes auratus</i>		
Western wood-pewee	<i>Contopus sordidulus</i>		
Say's phoebe	<i>Sayornis saya</i>		
Loggerhead shrike	<i>Lanius ludovicianus</i>		
Plumbeous vireo	<i>Vireo solitarius</i>		
Black-billed magpie	<i>Pica pica</i>		

Mammals

Cottontail	<i>Sylvilagus</i>
Eastern fox squirrel	<i>Sciurus niger</i>
American beaver	<i>Castor canadensis</i>
Western harvest mouse	<i>Reithrodontomys megalotis</i>
Deer mouse	<i>Peromyscus maniculatus</i>
Meadow vole	<i>Microtus pennsylvanicus</i>
Common muskrat	<i>Ondatra zibethicus</i>
Norway rat	<i>Rattus norvegicus*</i>
House mouse	<i>Mus musculus*</i>
Dog	<i>Canis familiaris*</i>
Coyote	<i>Canis latrans</i>
Red fox	<i>Vulpes vulpes</i>
Common raccoon	<i>Procyon lotor</i>
Cat	<i>Felis catus*</i>
Deer	<i>Odocoileus</i>

*Addresses of authors:***CHERIA A. JONES**

*Curator of Mammalogy
Department of Zoology
Denver Museum of Nature and Science
2001 Colorado Blvd.
Denver, CO 80210
cjones@dmns.org*

ELIZABETH A. DICKERSON

*Department of Zoology
Denver Museum of Nature and Science
2001 Colorado Blvd.
Denver, CO 80210
Beth.Dickerson@rf.doe.gov*


RONALD D. BEANE

*Research Associate
Department of Zoology
Denver Museum of Nature and Science
2001 Colorado Blvd.
Denver, CO 80210
rdbeane@email.msn.com*

*Current address:
U.S. Fish and Wildlife Service
Rocky Flats c/o Rocky Mountain
Arsenal National Wildlife Refuge
Building 121
Commerce City, CO 80022*

PUBLICATIONS OF THE MUSEUM OF TEXAS TECH UNIVERSITY

Institutional subscriptions are available through the Museum of Texas Tech University, attn: NSRL Publications Secretary, Box 43191, Lubbock, TX 79409-3191. Individuals may also purchase separate numbers of the Occasional Papers directly from the Museum of Texas Tech University.

**ISSN 0149-175X**

Museum of Texas Tech University, Lubbock, TX 79409-3191